

Physicians CARE®

Made in USA
2 Tablets

COLD & COUGH

<i>Active ingredients (in each tablet)</i>	<i>Purpose</i>
Acetaminophen 325mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 15mg	Cough suppressant
Guaifenesin 200mg.....	Expectorant
Phenylephrine HCl 5mg.....	Nasal decongestant

Uses

Temporarily relieves these cold symptoms

- cough
- sore throat
- minor aches and pains
- headaches
- nasal congestion
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

Temporarily reduces fever.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Store at 59°-86°F (15°-30°C) • Tamper-evident sealed packets • Do not use any opened or torn packets
Manufactured for Acme United Corp

Warnings (continued)

Do not use

- with any other drug containing acetaminophen
- with any other drug containing the above active ingredients
- if you are taking a prescription monoamine oxidase inhibitor (MAOI) drug

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- chronic cough that lasts as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase the drowsiness effect
- avoid alcoholic drinks
- be careful driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- new symptoms occur
- redness or swelling is present
- pain or nasal congestion gets worse or lasts for more than 7 days
- fever gets worse or lasts for more than 3 days
- you get nervous, dizzy or sleepless
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Adults and children: (12 years and older)	Take 2 tablets with water every 6-8 hours as needed. Do not take more than 8 tablets in 24 hours.
Children under 12 years:	Do not give to children under 12 years of age.

SEE CARTON FOR COMPLETE PRODUCT INFORMATION