

Joint Structure and Function

There are three types of equine joints: synovial, fibrous, and cartilaginous

Overview

The equine skeletal system is comprised of more than 200 bones that interconnect with the assistance of connective tissues such as tendons, ligaments, and cartilage.¹⁻³ Where two or more bones meet is considered a joint.

Of the three different types of joints, the synovial joint is the most common type in the horse's body. Synovial joints are freely movable, synovial fluid-filled anatomic structures that have a joint capsule surrounding the joint.^{1,3}

In contrast, a fibrous joint is an immovable joint that exists between two bones (such as the bones of the skull), and a cartilaginous joint holds bones together via fibrocartilaginous discs and ligaments that permit only a limited amount of movement. Examples of fibrous joints include the joints between individual vertebrae and the pubic symphysis (where the pubic bones join at the pelvic girdle).¹

The equine skeletal system is comprised of a number of different types of synovial joints. These include ball and socket joints in which the rounded end of one bone fits into the hollow curve of the opposing bone, such as the hip and shoulder; hinge joints in which the bones and ligaments are arranged to permit motion in one direction only, such as the elbow; and gliding joints, such as the carpus (knee) and tarsus (hock), in which the flat surfaces of opposing bones slide over one another, permitting only a limited amount of movement.

The Synovial (Diarthroidal) Joint

The basic structure of all synovial joints is the same, regardless of the type or location. All synovial joints have:⁴

- Two or more bones (ending with a plate of subchondral bone) covered with a thin layer of articular cartilage.
- A synovial fluid-filled cavity between the articulating bones.
- A synovial membrane lining the inner



The highly specialized tissues of the synovial joint (such as the hock) come together to perform two main functions: Enable movement and transfer load from one bone to another.

layer of the joint.

- A joint capsule that encapsulates the joint.

Depending on the joint, discs of cartilage (called menisci) might be present to cushion the articulating long bones inside the joint (such as the junction between the femur and tibia in the stifle). Also, the number and location of ligaments and tendons that help stabilize the joint varies from joint to joint. These include intra-articular ligaments (such as the cruciate ligaments in a horse's stifle) and ligaments and tendons located outside the joint (such as collateral ligaments).

Joints are Organs

A joint is more than simply the union of two or more bones; the joint is an organ.⁵ The highly specialized tissues of the synovial joint come together to perform two main functions: Enable movement and transfer load from one bone to another. In a normally functioning joint, both of these tasks are achieved in an efficient and pain-free manner. The secret of how this

frictionless, painless movement occurs relies on all the joint elements functioning in concert, but necessitates the involvement of healthy articular cartilage lining bones.

Articular cartilage is an extremely specialized connective tissue capable of withstanding very high loads during physical activity. It is composed largely of water (70-80% of articular cartilage is water), type II collagen fibrils, proteoglycan molecules such as aggrecan, chondrocytes (cartilage cells), and a variety of miscellaneous yet important molecules.


The chondrocytes are responsible for synthesizing, organizing, and regulating the extracellular matrix of the articular cartilage. The extracellular matrix is the tissue surrounding the chondrocytes where water, collagen, and proteoglycans are found. The type II collagen forms a fibrillar network within the extracellular matrix, which is responsible for maintaining the shape and strength of the tissue. Also found within the extracellular matrix are large, negatively-charged macromolecules called proteoglycans. These are a mixture of proteins

and long chains of sugar that attract large amounts of water, but repel each other. The most common proteoglycan in articular cartilage is the aggrecan—a very large proteoglycan that plays a pivotal role in the function of articular cartilage.

During weight bearing, the aggrecan molecules, which are already very tightly packed together, become further compressed. During this compression water molecules (that were attracted to the negatively-charged aggrecan molecules) are forced from the extracellular matrix of the cartilage, and all of the negatively charged branches of the aggrecan molecule repel each other like similar ends of a magnet. That is, the bones are protected by this layer of shock-absorbing articular cartilage, and the load is transmitted from one bone to another.

Joint Problems

Musculoskeletal dysfunction is the leading cause of lameness and loss of function in athletic horses. Joints can become damaged in one of two main ways: abnormal forces acting on normal cartilage or normal



View of normal left fetlock joint and abnormal (inflamed) right fetlock joint.

forces acting on abnormal cartilage.⁵

Horses also can have infections in various structures of the joint and congenital abnormalities (problems foals are born with, such as angular limb deformities) that affect joints. In a diseased joint, such as one with synovitis or capsulitis (i.e.,

inflammation of the synovial membrane or fibrous joint capsule, both common in athletic horses), even normal stresses can initiate, perpetuate, or exacerbate degeneration of articular cartilage.

The progressive destruction of articular cartilage in diarthroidal joints is known as osteoarthritis, which is the leading cause of joint disease in horses. 🐾

REFERENCES

1. Sellnow, L. Form and function of joints. www.TheHorse.com/36.
2. Sellnow, L. It all hinges on this. www.TheHorse.com/pdf/anatomy/anatomy2.pdf.
3. Loving, N.S. Equine skeletal system. www.TheHorse.com/15575.
4. Todhunter, R.J. Anatomy and physiology of synovial joints. McIlwraith, C.W.; Trotter, G.W., eds. *Joint Disease in the Horse*. Philadelphia: W.B. Saunders Company; 1-28.
5. McIlwraith, C.W. Osteoarthritis (degenerative joint disease): An update. *Proceedings of the 11th International Congress of the World Equine Veterinary Association*, Guarujá-SP, Brazil. 2009.

Further reading and free horse health e-newsletter: www.TheHorse.com/Joint-Problems

Authored by Stacey Oke, DVM, MSc; reviewed by Fairfield T. Bain, DVM, Dipl. ACVIM, ACVP, ACVECC.

DR. ROBIN PETERSON ILLUSTRATION



Joint Support Every Step of the Way



Platinum Performance® offers several joint support solutions so you can select the one that is right for your horse. Platinum Performance® CJ, a combination of three formulas: Platinum Performance® Equine, Ortho-Chon® II HA and ASU (Avocado/Soy Unsaponifiables) is the most comprehensive solution.

Recommended Platinum Performance® Products for Joint Health

Omega-3 FA	Vitamins, Trace Minerals	Antioxidants	Glucosamine	MSM, Cetyl Myristoleate, Boswelia	Hyaluronic Acid	ASU
Platinum Performance® Equine <i>Recommended For: Young & Sound Horses</i>						
Platinum Performance® Equine + Ortho-Chon® II <i>Recommended For: Senior Horses, Performance Horses</i>						
Platinum Performance® Equine + Ortho-Chon® HA <i>Recommended For: Senior Horses, Injured Horses</i>						
Platinum Performance® CJ <i>Recommended For: Performance Horses, Injured Horses</i>						

JOINT SUPPORT



Contact a Platinum Advisor to learn more about Platinum Performance® formulas that support Joint Health. Call 1-800-553-2400 or visit our website at www.PlatinumPerformance.com.

PLATINUM
PERFORMANCE®